

Marking the 50th Anniversary of *Immunology***Special regulatory T-cell review: A rose by any other name: from suppressor T cells to Tregs, approbation to unbridled enthusiasm**

Ronald N. Germain

Lymphocyte Biology Section, Laboratory of Immunology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA

doi:10.1111/j.1365-2567.2007.02779.x

Received 19 October 2007; accepted 1 November 2007.

Correspondence: Dr R. N. Germain, Lymphocyte Biology Section, Laboratory of Immunology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20892, USA. Email: rgermain@nih.gov

Summary

In the early 1970s a spate of papers by research groups around the world provided evidence for a negative regulatory role of thymus-derived lymphocytes (T cells). In 1971, Gershon and Kondo published a seminal paper in *Immunology* entitled 'Infectious Immunological Tolerance'¹ indicating that such negative regulation could be a dominant effect that prevented otherwise 'helpful' T cells from mediating their function. Over the next decade, suppressor T cells, as these negative regulatory cells became known, were intensively investigated and a complex set of interacting cells and soluble factors were described as mediators in this process of immune regulation. In the early 1980s, however, biochemical and molecular experiments raised questions about the interpretation of the earlier studies, and within a few years, the term 'suppressor T cell' had all but disappeared from prominence and research on this phenomenon was held in poor esteem. While this was happening, new studies appeared suggesting that a subset of T cells played a critical role in preventing autoimmunity. These T cells, eventually dubbed 'regulatory T cells', have become a major focus of modern cellular immunological investigation, with a predominance that perhaps eclipses even that seen in the earlier period of suppressor T cell ascendancy. This brief review summarizes the rise and fall of 'suppressorology' and the possibility that Tregs are a modern rediscovery of suppressor T cells made convincing by more robust models for their study and better reagents for their identification and analysis.

Keywords: immunoregulation; regulatory T cell; suppressor T cell; T lymphocyte**Introduction**

As much as we would all like to think otherwise, the conduct of science is as subject to human foibles such as ego, prejudice, and emotional-driven belief as many other societal activities. While we strive for a goal of dispassionate assessment of data, we cannot avoid coloration of our views by these factors nor avoid the impact of group-think. The history of studies on negative regulation of immune responses by subsets of T cells is a prime example. A large intellectual edifice involving suppressor T cells (Tsup) was established over a decade of intensive investigation by major laboratories, only to crumble with the publication of a limited number of studies that raised questions about certain reagents and molecular analyses. Many of those active in this area abandoned the field and the reputation of

those who persisted suffered greatly. Yet recent findings suggest that there was substantial truth to the basic phenomenology of T cell-mediated suppression and many of the observations made in these 'discredited' early analyses have been reprised in more modern studies of what are now called regulatory T cells (Tregs). Here I provide a brief overview of the rise and fall of Tsup and the reincarnation of these cells along with the theory of immunosuppression in the form of Tregs.

Early days

In the late 1960s evidence emerged that adaptive immunity was the product of two major classes of lymphocytes, B (bone-marrow derived) cells that made conventional immunoglobulin antibodies and T (thymus-derived) cells that were responsible for reactions such as delayed-type

hypersensitivity (DTH) and also involved in co-operation with B cells for generation of high-affinity antibody responses²⁻⁶. Within a short time, studies by several laboratories showed that T cells not only had these effector and positive co-operative roles, but also could depress immune responses^{1,7-16}. Early thoughts of mechanism revolved around B-cell destruction, but a seminal 1971 paper on infectious immunological tolerance by Gershon and Kondo¹, suggested that negative regulation could be mediated by interference with the activity of otherwise positive acting T cells. In concert with the emerging evidence for subsets of T cells, in particular as defined by anti-Ly antisera¹⁷⁻²⁵, this led to the notion that suppressor T cells were a specialized subset of lymphocytes whose role was to limit immune responses^{26,27}.

A confluence of rapidly emerging experimental evidence and the powerful logic of the argument that the immune system needed suitable brakes to prevent excessive activity led to rapid acceptance of suppressor T cells as a key paradigm of the growing field of cellular immunology. In short order, a number of experimental models involving proteins, red blood cells, haptens, and tumours (reviewed in refs²⁸⁻³⁸) were reported, all of which showed evidence of the activity of Tsup. Genes in the major histocompatibility complex (MHC), already known to control the positive aspects of immunity mediated by T cells, were found to also control suppressor T-cell function³⁹⁻⁴¹.

Research using a variety of antisera began to complicate the initial picture of Tsup function. An alloantiserum generated to a previously unrecognized subregion within the MHC, the so-called I-J subregion, was reported to react with Tsup and more strikingly, to be bind a soluble material derived from Tsup (TsF) that could substitute for intact cells in mediating negative immune regulation *in vitro* and *in vivo*⁴²⁻⁴⁴. Other antisera to specificities called Lyt1 and Lyt2 reacted in a differential manner with Tsup in different laboratories, a paradox whose solution was suggested to be the need for two distinct T-cell subsets to interact with each other for suppression to be manifest⁴⁵⁻⁴⁸.

The field became even more baroque under the influence of the contemporaneous paradigm of isotypic guidance of immune function⁴⁹. Anti-idiotypic antibodies made against immunoglobulins of a given specificity were claimed to recognize Tsup and TsF with a similar specificity and immunoglobulin H (IgH) allotypes were reported to provide another level of genetic restriction to the interactions involved in suppressive function⁵⁰⁻⁵⁶.

As these data on multigenic control emerged and other data were reported indicating that communication among the T-cell subsets involved in suppression could be mediated by a variety of distinct TsFs, some of which showed

both specificity for antigen and genetically restricted interactions, questions grew about whether the discordant nature of the complex models in the different experimental systems could be reconciled. Several attempts were made to provide an overarching theory of Tsup function (see ref. 34 for one such effort), most of which were based on the notion that different laboratories were interrogating distinct parts of a long concatenation of cell-cell interaction events that ended in the production of an antigen-unspecific effector factor that interfered with antigen-presenting cell function. However, larger problems loomed as molecular biology and monoclonal antibody technology began to bring a new rigour to the Wild West of immunological investigation of cell subsets and soluble mediators.

The beginning of the end

These complications involving many cell types and genetic restrictions notwithstanding, suppressor T cells remained an accepted part of the immunological firmament in the early 1980s. However, as molecular cloning began to have a major impact on biological studies in many fields, including immunology, and as the technique of Köhler and Milstein⁵⁷ began to yield monoclonal replacements for complex and not always reproducible or specific antisera, matters became problematic. Among many events that led to a loss of confidence in the Tsup story as it existed at this time, several stand out. One was the reported purification of an antigen-specific TsF⁵⁸, only to have another paper provide evidence that what had been identified was an apolipoprotein, not an antigen-specific T-cell derived molecule⁵⁹. Another was the analysis, first at the mRNA level⁶⁰ and then by genomic sequencing⁶¹, of the putative region in the MHC encoding I-J. No specific transcripts corresponding to this region were observed in Tsup and no nucleotide polymorphisms were found in DNA from the two strains of inbred mice [B10.A(3R) and B10.A(5R)] that were used to produce the initial alloantisera that identified I-J. An attempt was made to explain these data by postulating that I-J was actually an anti-idiotypic to the antigen-specific component of Ts and TsF⁶²⁻⁶⁴, which would result in the apparent mapping of I-J to the MHC because of the influence of such gene products on the combining site of the clonotypic receptors of T cells. However, this 'solution' of the I-J problem gained little traction. Finally, no specific T-cell receptor (TCR) β chain rearrangements were seen in DNA from putative Tsup hybridomas⁶⁵, although this result was less than definitive in that $\gamma\delta$ T cells also lacked such rearrangements in many cases.

Taken together, this spate of negative findings using the most modern tools for immunological investigation threw cold water on the entire field of Tsup and TsF. From a

field with great cache that dominated international meetings for years, it rapidly acquired a taint that impacted publication of papers in the area and perhaps even more significantly, the capacity of those heavily invested in the study of Tsup to maintain research funding. The result was a rapid loss of momentum and the development of the belief that many if not most of the observations reported in the area were flawed if not of dubious provenance. Citations for Tsup peaked as these events were unfolding and then rapidly declined to almost none.

The rise of Tregs

Like a phoenix, negative regulatory T cells rose from these ashes to a position of prominence in today's immunological thinking over precisely the interval from the demise of Tsup to the present. Insightful studies of autoimmunity arising in mice thymectomized early after birth suggested that a subset of T cells was critical for the restraint of effector development/function among more conventional T cells (reviewed in ref. 66). Close on the heels of these observations, other laboratories showed that transfer into immunodeficient hosts of purified T cells lacking activation/memory markers led to autoimmunity, especially inflammatory bowel disease, and that addition of a subset of T cells with certain memory cell markers could prevent this disease in a dominant manner^{67,68}. A report that inhibitory T cells were marked by high CD25 expression⁶⁹ and the demonstration that such cells limited TCR-induced T-cell proliferation of conventional CD4 and CD8 T cells in culture⁷⁰ opened up the field to study by the larger community, which rapidly confirmed that the memory/effector phenotype cells identified using CD45 and CD25 markers (Tregs) were important to preventing autoimmunity in lymphopenic animals given small numbers of conventional T cells and could also limit anti-pathogen immunity in certain cases (reviewed in refs 71–74).

The field made a major jump forward with the recognition that the transcription factor FoxP3 was critical for the development and/or function of these inhibitory Tregs^{75–77} and that either genetically modified reporter animals^{77–79} or monoclonal antibodies to FoxP3 could be used to identify these cells with some assurance. Perhaps most convincing that these Tregs played a major role in immune homeostasis was the dramatic autoimmune phenotype of mice with a mutation in the FoxP3 gene (scurfy mice) and the analogous autoimmune disease seen in humans with mutations in the FoxP3 gene (IPEX)^{80–83}.

Citations of papers describing Tregs is now rising in a manner akin to that seen in the early days of Tsup, and these days no broad-based immunological meeting worth its salt lacks a major session on these cells. The intense interest in this area of research is clearly evidenced by the

fact that such sessions are routinely oversubscribed, with attendees spilling out the doors of the lecture hall.

Are Tregs really Tsup in disguise?

There is little question that a distinct subset of T cells, most of which originate in the thymus, contribute to effective immune homeostasis. Whether all the claims made for a role of these so-called natural or nTregs in immune functioning will prove out in the long run is unclear, but their importance seems beyond question. Why have these cells been embraced so readily given the odious nature of Tsup, are the two types of inhibitory cells related, and what should the Treg field be careful about, given past history?

As to the first issue of acceptance, in contrast to the ephemeral nature of I–J, Tsup hybridomas, DTH measurements on mouse ears, and the like, Tregs are readily identified by widely available monoclonal reagents to CD25 and FoxP3, as well as by knock-in reporter mice expressing GFP in cells that transcribe the FoxP3 locus. Thus, many laboratories can study these cells with confidence that they are looking at more or less the same cell population (although some level of smug absoluteness is creeping into thinking on this issue and poses a serious risk going forward). Furthermore, using these tools, consistent data on the role of this subset in limiting autoimmune responses *in vivo* have emerged from a very large number of independent laboratories studying diverse model systems. In contrast to I–J on Tsup, there certainly is no question about the reality of CD25 and FoxP3 expression by cells that, as a population, can mediate negative immunoregulatory effects. For these reasons, the problems of the past that related to Tsup are not considered relevant to Tregs, at least in terms of their identity and reality.

With respect to whether Tregs and Tsup are related, this is a question that perhaps only those of us involved in the original Tsup work spend time considering. But it is instructive, I think, for a newer generation to appreciate that many of the observations made with regards to Tsup are strikingly similar to data relating to Tregs, and that even some of the most controversial of issues concerning Tsup are difficult to dismiss in light of modern knowledge and what was known (or more importantly, not known) when these early findings were reported.

1. An often reported characteristic of Tsup is that they could not be cloned *in vitro* using methods that reproducibly led to creation of lines and clones of conventional CD4 and CD8 T cells; this property is entirely consistent with the well-accepted anergic character of Tregs *in vitro*^{71,73,74} and the difficulty of driving their proliferation in culture except with heroic concentra-

- tions of interleukin-2 (IL-2) and strong stimulation with anti-TCR and CD28 antibodies.⁸⁴
2. The *in vivo* function of Tsup was eliminated by treatment with low dose cyclophosphamide⁸⁵; this has also been reported to be true for Treg function.⁸⁶
 3. Natural Tregs are CD4 T cells that are selected in the thymus by recognition of MHC class II molecules^{71,73,74,77} and that function in the periphery as do other CD4 T cells, using MHC class II molecules for antigen recognition; likewise, the genes regulating Tsup function were mapped to the class II region of the MHC.^{40,41}
 4. The antigen recognition unit of T cells is a disulfide-bonded heterodimer with two chains in the ~40–50 000 MW range⁸⁷; reports on the molecular nature of the antigen-specific suppressor factor of Tsup characterized the material as a disulfide-linked heterodimer of similar molecular mass^{88,89}; it would be quite fortuitous for the authors of the latter work to have arrived at this result by chance, given that only immunoglobulins of much greater molecular mass were known to be antigen-specific molecules at the time.
 5. The lack of TCR β rearrangements in Tsup hybridomas involved studies of DNA from long-term cultures of these notoriously unstable cells, without repeated selection for antigen-specificity and without the possibility of sorting for expressed TCR because the relevant antibodies were not available; selection of CD3⁺ cells from such cultures once the proper reagents became available showed that these cells did express conventional TCR and that increasing the proportion of TCR⁺ cells from a few percent to close to homogeneity also increased suppressive activity by a comparable extent⁹⁰; likewise, authentic TCR $\alpha\beta$ determinants were found on TsF from Tsup when the proper monoclonal reagents became available.⁹¹
 6. The ability of Tsup and TsF to bind antigen in the absence of MHC class II molecules is seemingly problematic given our knowledge of T-cell receptor structure-function and recognition of peptide-MHC molecule ligands; however, several reports have shown that T cells with a functional requirement for antigen presentation by MHC class I or II molecules have TCR that, when isolated biochemically, can show direct binding to certain antigens independent of MHC molecules, as claimed for Tsup and TsF.^{92–94}
 7. Infectious tolerance has been rediscovered^{95,96} and is now an increasingly popular view of how Tregs work; although the mediator of this infectious process is considered to be an antigen-unspecific cytokine, the idea that the potency of suppression is amplified by recruitment of new cells into the suppressive pool parallels the Ts1-Ts2-Ts3 schemes from the old Tsup days, as does the evidence that the ultimate mediation of Treg activity involves any of several immunosuppressive

cytokines (transforming growth factor- β (TGF- β) and IL-10 chief among these^{71,73,74,97}) that exert their activity via antigen presenting cells, just as concluded in the early Tsup studies.⁹⁸

Finally, what lessons should be drawn from the rise and fall of Tsup and the recent impact of Treg studies? At a minimum, given the list above, it seems that the field threw the baby out with the bathwater. Many of the biological observations made in connection with Tsup seem quite robust and similar if not identical to those now being found true of Tregs. The major problem in the earlier work was the inability of those involved to move the field forward on the molecular level and the possibility that some key observations were misleading, if not just plain wrong. But do any of us imagine that all the aspects of the biology of Tregs that are well accepted today are absolutely correct? Already questions are being raised about whether the widely used *in vitro* assay for suppression of proliferation is related more to competition for cytokines than to the better accepted mechanisms mediating the *in vivo* physiological function of these cells⁹⁹. Likewise, the cell-cell contact requirement for suppression claimed in the culture assay may not be what it seems in terms of real Treg function^{100,101}. Early reports of Treg function being entirely divorced from TGF- β activity^{102,103} are being reconsidered in light of strong data to the contrary (reviewed in ref. 97), but this change in viewpoint has not resulted in a push to discredit all the work on Tregs done prior this point in time. The absolute linkage of FoxP3 expression with Tregs has been called into question by data from human cells, which express this protein without necessarily showing the *trans*-regulatory properties of mouse Tregs¹⁰⁴. Yet the field does not question the essential role of FoxP3 in Treg development and function. Finally, even in the mouse, a debate now rages about the occurrence and relevance of induced Tregs¹⁰⁵, but this is a more typical scientific dispute within a field that accepts the underlying existence of regulatory cells expressing FoxP3.

In other words, the fact that errors of interpretation and inadequate experiments exist in the Treg arena, just as they did in the Tsup era, has not done in the field in the way that a few reports disputing aspects of the 'received wisdom' in the Tsup field did 25 years ago. Perhaps we have all grown up a bit since then and have a more sophisticated view of the complexity of the immune system and our limited capacity to get everything right in every report probing its workings. Possibly, the field has also learned a lesson from the self-inflicted wound of the early 1980s when the results of serious scientists were discredited without adequate consideration of alternative interpretations for the experimental data, and without seeking to separate the wheat from the chaff.

Indeed, if more effort gone into separating the widely reproduced and fundamental aspects of Tsup function

from less certain claims introduced by limited technology and suboptimal experiments, we might be even further along than we are in understanding immune regulation. At the very least, immunology would not have given itself such a black-eye in the view of other biologists and we would have less explaining to do about whether we pursue our work with the rigor of other fields. Some of the same internal derogation of other investigators in the field is beginning to creep into the Treg arena – it seems better for us all to carefully question potentially incorrect conclusions on their merits and use our concern about these matters to drive better experiments, than to engage in the type of self-defeating internecine warfare that led to the demise of the Tsup field in the past.

References

- 1 Gershon RK, Kondo K. Infectious immunological tolerance. *Immunology* 1971; **21**:903–14.
- 2 Miller JF. Immunological function of the thymus. *Lancet* 1961; **2**:748–9.
- 3 Good RA, Dalmasso AP, Martinez C, Archer OK, Pierce JC, Papermaster BW. The role of the thymus in development of immunologic capacity in rabbits and mice. *J Exp Med* 1962; **116**:733–96.
- 4 Claman HN, Chaperon EA, Triplett RF. Thymus-marrow cell combinations. Synergism in antibody production. *Proc Soc Exp Biol Med* 1966; **122**:1167–71.
- 5 Mosier DE. A requirement for two cell types for antibody formation *in vitro*. *Science* 1967; **158**:1573–5.
- 6 Mitchell GF, Miller JF. Cell to cell interaction in the immune response. II. The source of hemolysin-forming cells in irradiated mice given bone marrow and thymus or thoracic duct lymphocytes. *J Exp Med* 1968; **128**:821–37.
- 7 Baker PJ, Stashak PW, Amsbaugh DF, Prescott B, Barth RF. Evidence for the existence of two functionally distinct types of cells which regulate the antibody response to type 3 pneumococcal polysaccharide. *J Immunol* 1970; **105**:1581–3.
- 8 Okumura K, Tada T. Regulation of homocytotropic antibody formation in the rat. VI. Inhibitory effect of thymocytes on the homocytotropic antibody response. *J Immunol* 1971; **107**:1682–9.
- 9 Droege W. Amplifying and suppressive effect of thymus cells. *Nature* 1971; **234**:549–51.
- 10 Allison AC, Denman AM, Barnes RD. Cooperating and controlling functions of thymus-derived lymphocytes in relation to autoimmunity. *Lancet* 1971; **2**:135–40.
- 11 Jacobson EB, Herzenberg LA. Active suppression of immunoglobulin allotype synthesis. I. Chronic suppression after perinatal exposure to maternal antibody to paternal allotype in (SJL × BALB-c)F1 mice. *J Exp Med* 1972; **135**:1151–62.
- 12 Kerbel RS, Eidinger D. Enhanced immune responsiveness to a thymus-independent antigen early after adult thymectomy: evidence for short-lived inhibitory thymus-derived cells. *Eur J Immunol* 1972; **2**:114–8.
- 13 Yoshinaga M, Yoshinaga A, Waksman BH. Regulation of lymphocyte responses *in vitro*. I. Regulatory effect of macrophages and thymus-dependent (T) cells on the response of thymus-independent (B) lymphocytes to endotoxin. *J Exp Med* 1972; **136**:956–61.
- 14 Rich RR, Pierce CW. Biological expressions of lymphocyte activation. II. Generation of a population of thymus-derived suppressor lymphocytes. *J Exp Med* 1973; **137**:649–59.
- 15 Katz DH, Paul WE, Benacerraf B. Carrier function in anti-hapten antibody responses. VI. Establishment of experimental conditions for either inhibitory or enhancing influences of carrier-specific cells on antibody production. *J Immunol* 1973; **110**:107–17.
- 16 Okumura K, Tada T. Suppression of hapten-specific antibody response by carrier-specific T cells. *Nat New Biol* 1973; **245**:180–2.
- 17 Cantor H, Asofsky R. Synergy among lymphoid cells mediating the graft-versus-host response. II. Synergy in graft-versus-host reactions produced by BALB-c lymphoid cells of differing anatomic origin. *J Exp Med* 1970; **131**:235–46.
- 18 Cantor H, Asofsky R. Synergy among lymphoid cells mediating the graft-versus-host response. 3. Evidence for interaction between two types of thymus-derived cells. *J Exp Med* 1972; **135**:764–79.
- 19 Schendel DJ, Alter BJ, Bach FH. The involvement of LD- and SD-region differences in MLC and CML: a three-cell experiment. *Transplant Proc* 1973; **5**:1651–5.
- 20 Dennert G. Evidence for non-identity of T killer and T helper cells sensitised to allogeneic cell antigens. *Nature* 1974; **249**:358–60.
- 21 Tigelaar RE, Gorczynski RM. Separable populations of activated thymus-derived lymphocytes identified in two assays for cell-mediated immunity to murine tumor allografts. *J Exp Med* 1974; **140**:267–89.
- 22 Shiku H, Kisielow P, Bean MA, Takahashi T, Boyse EA, Oettgen HF, Old LJ. Expression of T-cell differentiation antigens on effector cells in cell-mediated cytotoxicity *in vitro*. Evidence for functional heterogeneity related to the surface phenotype of T cells. *J Exp Med* 1975; **141**:227–41.
- 23 Cantor H, Boyse EA. Functional subclasses of T-lymphocytes bearing different Ly antigens. I. The generation of functionally distinct T-cell subclasses is a differentiative process independent of antigen. *J Exp Med* 1975; **141**:1376–89.
- 24 Cantor H, Boyse EA. Functional subclasses of T lymphocytes bearing different Ly antigens. II. Cooperation between subclasses of Ly+ cells in the generation of killer activity. *J Exp Med* 1975; **141**:1390–9.
- 25 Cantor H, Shen FW, Boyse EA. Separation of helper T cells from suppressor T cells expressing different Ly components. II. Activation by antigen: after immunization, antigen-specific suppressor and helper activities are mediated by distinct T-cell subclasses. *J Exp Med* 1976; **143**:1391–40.
- 26 Vadas MA, Miller JF, McKenzie IF, Chism SE, Shen FW, Boyse EA, Gamble JR, Whitelaw AM. Ly and Ia antigen phenotypes of T cells involved in delayed-type hypersensitivity and in suppression. *J Exp Med* 1976; **144**:10–9.
- 27 Okumura K, Takemori T, Tokuhisa T, Tada T. Specific enrichment of the suppressor T cell bearing I-J determinants: parallel functional and serological characterizations. *J Exp Med* 1977; **146**:1234–45.
- 28 Tada T, Taniguchi M, Takemori T. Properties of primed suppressor T cells and their products. *Transplant Rev* 1975; **26**:106–29.

- 29 Herzenberg LA, Okumura K, Metzler CM. Regulation of immunoglobulin and antibody production by allotype suppressor T cells in mice. *Transplant Rev* 1975; **27**:57–83.
- 30 Kapp JA, Pierce CW, Theze J, Benacerraf B. Modulation of immune responses by suppressor T cells. *Fed Proc* 1978; **37**:2361–4.
- 31 Claman HN, Miller SD, Sy MS, Moorhead JW. Suppressive mechanisms involving sensitization and tolerance in contact allergy. *Immunol Rev* 1980; **50**:105–32.
- 32 Greene MI, Perry LL, Benacerraf B. Regulation of the immune response to tumor antigen. *Am J Pathol* 1979; **95**:159–69.
- 33 Feldmann M, Kontiainen S. Antigen specific T cell factors. *Mol Cell Biochem* 1980; **30**:177–93.
- 34 Benacerraf B, Germain RN. A single major pathway of T-lymphocyte interactions in antigen-specific immune suppression. *Scand J Immunol* 1981; **13**:1–10.
- 35 Greene MI, Nelles MJ, Sy MS, Nisonoff A. Regulation of immunity to the azobenzenearsonate hapten. *Adv Immunol* 1982; **32**:253–300.
- 36 Dorf ME, Benacerraf B. Suppressor cells and immunoregulation. *Annu Rev Immunol* 1984; **2**:127–57.
- 37 Asherson GL, Colizzi V, Zembala M. An overview of T-suppressor cell circuits. *Annu Rev Immunol* 1986; **4**:37–68.
- 38 Green DR, Flood PM, Gershon RK. Immunoregulatory T-cell pathways. *Annu Rev Immunol* 1983; **1**:439–63.
- 39 Kapp JA, Pierce CW, Schlossman S, Benacerraf B. Genetic control of immune responses *in vitro*. V. Stimulation of suppressor T cells in nonresponder mice by the terpolymer L-glutamic acid 60-L-alanine 30-L-tyrosine 10 (GAT). *J Exp Med* 1974; **140**:648–59.
- 40 Debre P, Kapp JA, Benacerraf B. Genetic control of specific immune suppression. I. Experimental conditions for the stimulation of suppressor cells by the copolymer L-glutamic acid50-L-tyrosine50 (GT) in nonresponder BALB/c mice. *J Exp Med* 1975; **142**:1436–46.
- 41 Debre P, Kapp JA, Dorf ME, Benacerraf B. Genetic control of specific immune suppression. II. H-2-linked dominant genetic control of immune suppression by the random copolymer L-glutamic acid50-L-tyrosine50 (GT). *J Exp Med* 1975; **142**:1447–54.
- 42 Murphy DB, Herzenberg LA, Okumura K, McDevitt HO. A new I subregion (I-J) marked by a locus (Ia-4) controlling surface determinants on suppressor T lymphocytes. *J Exp Med* 1976; **144**:699–712.
- 43 Okumura K, Herzenberg LA, Murphy DB, McDevitt HO. Selective expression of H-2 (i-region) loci controlling determinants on helper and suppressor T lymphocytes. *J Exp Med* 1976; **144**:685–98.
- 44 Tada T, Taniguchi M, David CS. Properties of the antigen-specific suppressive T-cell factor in the regulation of antibody response of the mouse. IV. Special subregion assignment of the gene(s) that codes for the suppressive T-cell factor in the H-2 histocompatibility complex. *J Exp Med* 1976; **144**:713–25.
- 45 Eardley DD, Hugenberger J, McVay-Boudreau L, Shen FW, Gershon RK, Cantor H. Immunoregulatory circuits among T-cell sets. I. T-helper cells induce other T-cell sets to exert feedback inhibition. *J Exp Med* 1978; **147**:1106–15.
- 46 Cantor H, Hugenberger J, McVay-Boudreau L, Eardley DD, Kemp J, Shen FW, Gershon RK. Immunoregulatory circuits among T-cell sets. Identification of a subpopulation of T-helper cells that induces feedback inhibition. *J Exp Med* 1978; **148**:871–7.
- 47 McDougal JS, Shen FW, Elster P. Generation of T helper cells *in vitro*. V. Antigen-specific Ly1⁺ T cells mediate the helper effect and induce feedback suppression. *J Immunol* 1979; **122**:437–42.
- 48 McDougal JS, Shen FW, Cort SP, Bard J. Feedback suppression: phenotypes of T cell subsets involved in the Ly1 T cell-induced immunoregulatory circuit. *J Immunol* 1980; **125**:1157–60.
- 49 Jerne NK. Idiotypic networks and other preconceived ideas. *Immunol Rev* 1984; **79**:5–24.
- 50 Germain RN, Ju ST, Kipps TJ, Benacerraf B, Dorf ME. Shared idiotypic determinants on antibodies and T-cell-derived suppressor factor specific for the random terpolymer L-glutamic acid60-L-alanine30-L-tyrosine10. *J Exp Med* 1979; **149**:613–22.
- 51 Bach BA, Greene MI, Benacerraf B, Nisonoff A. Mechanisms of regulation of cell-mediated immunity. IV. Azobenzenearsonate-specific suppressor factor(s) bear cross-reactive idiotypic determinants the expression of which is linked to the heavy-chain allotype linkage group of genes. *J Exp Med* 1979; **149**:1084–98.
- 52 Eardley DD, Shen FW, Cantor H, Gershon RK. Genetic control of immunoregulatory circuits. Genes linked to the Ig locus govern communication between regulatory T-cell sets. *J Exp Med* 1979; **150**:44–50.
- 53 Weinberger JZ, Germain RN, Ju ST, Greene MI, Benacerraf B, Dorf ME. Hapten-specific T-cell responses to 4-hydroxy-3-nitrophenyl acetyl. II. Demonstration of idiotypic determinants on suppressor T cells. *J Exp Med* 1979; **150**:761–76.
- 54 Sy MS, Dietz MH, Germain RN, Benacerraf B, Greene MI. Antigen- and receptor-driven regulatory mechanisms. IV. Idiotype-bearing I-J⁺ suppressor T cell factors induce second-order suppressor T cells which express anti-idiotypic receptors. *J Exp Med* 1980; **151**:1183–95.
- 55 Weinberger JZ, Benacerraf B, Dorf ME. Hapten-specific T cell responses to 4-hydroxy-3-nitrophenyl acetyl. III. Interaction of effector suppressor T cells is restricted by H-2 and Igh-V genes. *J Exp Med* 1980; **151**:1413–23.
- 56 Takei I, Sumida T, Taniguchi M. Acceptor-suppressor T cell hybridoma with a receptor recognizing antigen-specific suppressor factor. *J Exp Med* 1983; **158**:1912–23.
- 57 Kohler G, Milstein C. Continuous cultures of fused cells secreting antibody of predefined specificity. *Nature* 1975; **256**:495–7.
- 58 Krupen K, Araneo BA, Brink L, Kapp JA, Stein S, Wieder KJ, Webb DR. Purification and characterization of a monoclonal T-cell suppressor factor specific for poly(LGlu60LAla30LTyr10). *Proc Natl Acad Sci U S A* 1982; **79**:1254–8.
- 59 Breslow JL, Ross D, McPherson J, Williams H, Kurnit D, Nussbaum AL, Karathanasis SK, Zannis VI. Isolation and characterization of cDNA clones for human apolipoprotein A-I. *Proc Natl Acad Sci U S A* 1982; **79**:6861–5.
- 60 Kronenberg M, Steinmetz M, Kober J *et al*. RNA transcripts for I-J polypeptides are apparently not encoded between the I-A and I-E subregions of the murine major histocompatibility complex. *Proc Natl Acad Sci U S A* 1983; **80**:5704–8.
- 61 Kober JA, Strauss E, Minard K, Hood L. Molecular analysis of the hotspot of recombination in the murine major histocompatibility complex. *Science* 1986; **234**:173–9.
- 62 Schrader JW. Nature of the T-cell receptor. Both the T-cell receptor and antigen-specific T-cell-derived factors are coded for by V genes but express anti-self idiotypes indirectly determined

- by major histocompatibility complex genes. *Scand J Immunol* 1979; **10**:387–93.
- 63 Uracz W, Abe R, Tada T. Involvement of I-J epitopes in the self- and allo-recognition sites of T cells: blocking of syngeneic and allogeneic mixed lymphocyte reaction-responder cells by monoclonal anti-I-J antibodies. *Proc Natl Acad Sci U S A* 1985; **82**:2905–9.
 - 64 Sumida T, Sado T, Kojima M, Ono K, Kamisaku H, Taniguchi M. I-J as an idiotype of the recognition component of antigen-specific suppressor T-cell factor. *Nature* 1985; **316**:738–41.
 - 65 Hedrick SM, Germain RN, Bevan MJ *et al.* Rearrangement and transcription of a T-cell receptor beta-chain gene in different T-cell subsets. *Proc Natl Acad Sci U S A* 1985; **82**:531–5.
 - 66 Nishizuka Y. A novel experimental system of organ-localized autoimmune diseases in the mouse. *Acta Pathol Jpn* 1982; **32**(Suppl. 1):211–22.
 - 67 Powrie F, Mason D. OX-22high CD4⁺ T cells induce wasting disease with multiple organ pathology: prevention by the OX-22low subset. *J Exp Med* 1990; **172**:1701–8.
 - 68 Powrie F, Leach MW, Mauze S, Caddle LB, Coffman RL. Phenotypically distinct subsets of CD4⁺ T cells induce or protect from chronic intestinal inflammation in C. B-17 scid mice. *Int Immunol* 1993; **5**:1461–71.
 - 69 Sakaguchi S, Sakaguchi N, Asano M, Itoh M, Toda M. Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor alpha-chains (CD25). Breakdown of a single mechanism of self-tolerance causes various autoimmune diseases. *J Immunol* 1995; **155**:1151–64.
 - 70 Thornton AM, Shevach EM. CD4⁺ CD25⁺ immunoregulatory T cells suppress polyclonal T cell activation *in vitro* by inhibiting interleukin 2 production. *J Exp Med* 1998; **188**:287–96.
 - 71 Shevach EM. Regulatory T cells in autoimmunity. *Annu Rev Immunol* 2000; **18**:423–49.
 - 72 Gavin M, Rudensky A. Control of immune homeostasis by naturally arising regulatory CD4⁺ T cells. *Curr Opin Immunol* 2003; **15**:690–6.
 - 73 Sakaguchi S, Sakaguchi N. Regulatory T cells in immunologic self-tolerance and autoimmune disease. *Int Rev Immunol* 2005; **24**:211–26.
 - 74 Banham AH, Powrie FM, Suri-Payer E. FOXP3⁺ regulatory T cells: current controversies and future perspectives. *Eur J Immunol* 2006; **36**:2832–6.
 - 75 Hori S, Nomura T, Sakaguchi S. Control of regulatory T cell development by the transcription factor Foxp3. *Science* 2003; **299**:1057–61.
 - 76 Fontenot JD, Gavin MA, Rudensky AY. Foxp3 programs the development and function of CD4⁺ CD25⁺ regulatory T cells. *Nat Immunol* 2003; **4**:330–6.
 - 77 Fontenot JD, Rasmussen JP, Williams LM, Dooley JL, Farr AG, Rudensky AY. Regulatory T cell lineage specification by the forkhead transcription factor foxp3. *Immunity* 2005; **22**:329–41.
 - 78 Wan YY, Flavell RA. Identifying Foxp3-expressing suppressor T cells with a bicistronic reporter. *Proc Natl Acad Sci U S A* 2005; **102**:5126–31.
 - 79 Bettelli E, Carrier Y, Gao W, Korn T, Strom TB, Oukka M, Weiner HL, Kuchroo VK. Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells. *Nature* 2006; **441**:235–8.
 - 80 Wildin RS, Ramsdell F, Peake J *et al.* X-linked neonatal diabetes mellitus, enteropathy and endocrinopathy syndrome is the human equivalent of mouse scurfy. *Nat Genet* 2001; **27**:18–20.
 - 81 Bennett CL, Christie J, Ramsdell F *et al.* The immune dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome (IPEX) is caused by mutations of FOXP3. *Nat Genet* 2001; **27**:20–1.
 - 82 Brunkow ME, Jeffery EW, Hjerrild KA *et al.* Disruption of a new forkhead/winged-helix protein, scurfy, results in the fatal lymphoproliferative disorder of the scurfy mouse. *Nat Genet* 2001; **27**:68–73.
 - 83 Khattry R, Cox T, Yasayko SA, Ramsdell F. An essential role for Scurfin in CD4⁺ CD25⁺ T regulatory cells. *Nat Immunol* 2003; **4**:337–42.
 - 84 Tang Q, Henriksen KJ, Bi M *et al.* In vitro-expanded antigen-specific regulatory T cells suppress autoimmune diabetes. *J Exp Med* 2004; **199**:1455–65.
 - 85 Polak L, Turk JL. Reversal of immunological tolerance by cyclophosphamide through inhibition of suppressor cell activity. *Nature* 1974; **249**:654–6.
 - 86 Ghiringhelli F, Larmonier N, Schmitt E *et al.* CD4⁺ CD25⁺ regulatory T cells suppress tumor immunity but are sensitive to cyclophosphamide which allows immunotherapy of established tumors to be curative. *Eur J Immunol* 2004; **34**:336–44.
 - 87 Allison JP, Lanier LL. Identification of antigen receptor-associated structures on murine T cells. *Nature* 1985; **314**:107–9.
 - 88 Saito T, Taniguchi M. Chemical features of an antigen-specific suppressor T cell factor composed of two polypeptide chains. *J Mol Cell Immunol* 1984; **1**:137–45.
 - 89 Taniguchi M, Saito T, Takei I, Tokuhisa T. Presence of inter-chain disulfide bonds between two gene products that compose the secreted form of an antigen-specific suppressor factor. *J Exp Med* 1981; **153**:1672–7.
 - 90 Kuchroo VK, Steele JK, Billings PR, Selvaraj P, Dorf ME. Expression of CD3-associated antigen-binding receptors on suppressor T cells. *Proc Natl Acad Sci U S A* 1988; **85**:9209–13.
 - 91 Fairchild RL, Kubo RT, Moorhead JW. Soluble factors in tolerance and contact sensitivity to 2,4-dinitro-fluorobenzene in mice. IX. A monoclonal T cell suppressor molecule is structurally and serologically related to the alpha/beta T cell receptor. *J Immunol* 1988; **141**:3342–8.
 - 92 Siliciano RF, Hemesath TJ, Pratt JC, Dintzis RZ, Dintzis HM, Acuto O, Shin HS, Reinherz EL. Direct evidence for the existence of nominal antigen binding sites on T cell surface T_H1 alpha-beta heterodimers of MHC-restricted T cell clones. *Cell* 1986; **47**:161–71.
 - 93 Kuchroo VK, Steele JK, O'Hara RM Jr, Jayaraman S, Selvaraj P, Greenfield E, Kubo RT, Dorf ME. Relationships between antigen-specific helper and inducer suppressor T cell hybridomas. *J Immunol* 1990; **145**:438–48.
 - 94 Diamond DJ, Szalay P, Symer D *et al.* Major histocompatibility complex independent T cell receptor-antigen interaction: functional analysis using fluorescein derivatives. *J Exp Med* 1991; **174**:229–41.
 - 95 Waldmann H, Adams E, Fairchild P, Cobbold S. Infectious tolerance and the long-term acceptance of transplanted tissue. *Immunol Rev* 2006; **212**:301–13.
 - 96 Qin S, Cobbold SP, Pope H, Elliott J, Kioussis D, Davies J, Waldmann H. 'Infectious' transplantation tolerance. *Science* 1993; **259**:974–7.

- 97 Wan YY, Flavell RA. Regulatory T cells, transforming growth factor-beta, and immune suppression. *Proc Am Thorac Soc* 2007; **4**:271–6.
- 98 Ptak W, Zembala M, Asherson GL, Marcinkiewicz J. Inhibition of contact sensitivity by macrophages. *Int Arch Allergy Appl Immunol* 1981; **65**:121–8.
- 99 Pandiyan P, Zheng L, Ishihara S, Reed J, Lenardo MJ. CD4⁺ CD25⁺ Foxp3⁺ regulatory T cells induce cytokine deprivation-mediated apoptosis of effector CD4⁺ T cells. *Nat Immunol* 2007; **8**:1353–62.
- 100 Thornton AM, Shevach EM. Suppressor effector function of CD4⁺ CD25⁺ immunoregulatory T cells is antigen nonspecific. *J Immunol* 2000; **164**:183–90.
- 101 Mempel TR, Pittet MJ, Khazaie K, Weninger W, Weissleder R, von Boehmer H, von Andrian UH. Regulatory T cells reversibly suppress cytotoxic T cell function independent of effector differentiation. *Immunity* 2006; **25**:129–41.
- 102 Piccirillo CA, Letterio JJ, Thornton AM, McHugh RS, Mamura M, Mizuhara H, Shevach EM. CD4(+)CD25(+) regulatory T cells can mediate suppressor function in the absence of transforming growth factor beta1 production and responsiveness. *J Exp Med* 2002; **196**:237–46.
- 103 Kullberg MC, Hay V, Cheever AW, Mamura M, Sher A, Letterio JJ, Shevach EM, Piccirillo CA. TGF-beta1 production by CD4⁺ CD25⁺ regulatory T cells is not essential for suppression of intestinal inflammation. *Eur J Immunol* 2005; **35**:2886–95.
- 104 Gavin MA, Torgerson TR, Houston E *et al*. Single-cell analysis of normal and FOXP3-mutant human T cells: FOXP3 expression without regulatory T cell development. *Proc Natl Acad Sci USA* 2006; **103**:6659–64.
- 105 Wong J, Mathis D, Benoist CT. CR-based lineage tracing: no evidence for conversion of conventional into regulatory T cells in response to a natural self-antigen in pancreatic islets. *J Exp Med* 2007; **204**:2039–45.

